

Effect of total pelvic peritonectomy*(including DIE lesion) in cases of Deep infiltrative endometriosis(DIE) with intractable pain

Abstract ID : 2350

Submitted by : dogyun kim the 2016-12-24 08:10:17

Category : SEUD CONGRESS

Typology : Video

Status : Validated

Authorisation to disclose : Yes/Oui

We introduce our laparoscopic pelvic peritonectomy technique including removal of Deep Infiltrative Endometriosis(DIE) lesion and results and effect of it. The 150 DIE patients with intractable pain had treated by laparoscopic pelvic peritonectomy with DIE lesion removal at Pohang Saint Mary hospital, South Korea from 1st day. January, 2013 to last day. December. 2014.

2 years later, 132 patients answer our question paper and check by transvaginal sonographic test. The VAS score had changed from 8.5 to 2.5.

The resolving of intractable pain was very effective. The recurrence rate of DIE was about 15%.

The our pelvic peritonectomy with removal of all DIE lesion(uterosacral ligament, rectovaginal septum, rectum, bladder et al) is very effective surgical choice in cases of DIE with intractable pain.

Keywords : pelvic peritonectomy, Deep infiltrative endometriosis

Authors :

Files are accepted up to a size of:

[1482561747_mis-.wmv](#)

[1482561747_pelvic-peritonectomy.wmv](#)

References : , , ,

Authors

Dogyun Kim 1,

1. Gynecology, Pohang St Mary hospital, Pohang, KOREA, REPUBLIC OF

Authors (raw format)

Kim Dogyun - email : gim555@hanmail.net Institution : Pohang St Mary hospital Department : Gynecology City : Pohang Country : KOREA, REPUBLIC OF Speaker : Yes

